

Med2000

Succorol[®]

Succorol[®] Patches, 200 mg

Compare to the Tylenol^{®†} tablets active ingredient!

Pain Reliever / Guilt Reducer

(NCCCD)

30 tablets

▣ ≤ actual size

**TAMPER EVIDENT — DO NOT USE IF CARTON IS OPENED OR SAFETY SEAL
UNDER CAP IS BROKEN OR MISSING**

Drug Facts

Active ingredient (in each tablet) and purpose:

- Succorol[®]pain reliever
- 200 mg (NCCCD)*guilt reducer

*non-consensual cultural culpability drug

Uses

patches aid both recipient and administrator. temporarily relieve pains and blames due to:

- headache

- heartache
- backache
- muscular aches
- menstrual cramps
- parenthood
- alcoholism
- heavy hands
- destitution
- stretch marks
- postpartum depression
- loss or absence
- whitewashing / cultural betrayal
- abandonment
- theft
- sacrifice
- dreams
- obligation
- uppity-ness
- patriarchy
- childlessness
- fatigue
- hair loss
- anger management issues

- empty apologies
- nightmares
- autoimmune disease
- swelling
- cancer

Warnings

Allergy alert: Succorol[®] may cause a severe allergic reaction, especially in hypocrites or people allergic to conscience.

Symptoms may include:

- shock
- asthma (wheezing)
- melting
- hallucinations
- vivid reminiscences
- phantom sensory stimulation, especially auditory
- cognitive dissonance
- out-of-body sensations

Stomach bleeding warning: This product contains a non-consensual cultural culpability drug (NCCCD), which may trigger severe stomach bleeding via internal karmic mechanism. The chance is higher if you:

- are age 60 or older
- have a dark complexion

- have had stress ulcers or anger problems
- take a blood thinning (anticoagulant) or pain relieving drug [morphine, ibuprofen, or others]
- have significant non-Anglo-Saxon ancestry
- lack immune resistance to other drugs containing prescription or nonprescription NCCCDs
- have 3 or more alcoholic drinks every day while using this product
- suffer from narcissistic, obdurate, and/or sadistic tendencies
- take a higher dose or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NCCCDs increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/guilt reducer
- right before or after radiotherapy

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or guilt reducers
- you have a history of
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

- you are taking a diuretic or antibiotics

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food, milk, or intimacy if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following symptoms:

- feel faint
- feel remorseful
- feel remorseless
- vomit blood
- have pain that does not get better
- chest pain
- trouble breathing
- weakness in one part or side of the body or personality
- redness or swelling is present in painful areas
- any new symptoms appear

Drug Facts (continued)

Side Effects

extended and/or incorrect use may induce:

- backache

- muscular aches
- cognitive dissonance
- resentment
- carpal tunnel syndrome
- dry skin and rash
- intrusive or violent thoughts
- impulsive behaviour
- migraine
- rebellion
- heart palpitations
- reclamation of power
- reclamation of agency
- alcoholism
- projection
- anger displacement
- cyclic psychological tendencies
- maladaptive coping mechanisms
- heartache
- suicidal thoughts
- general crisis

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use Succorol[®] in excess during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may result in perinatal transmission of NCCC substances.

Keep out of reach of children and wifes. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults: take 1 patch every 4 to 6 hours while blames persist
 - if pains do not respond to 1 patch, 2 patches may be used
 - do not exceed 3 patches in 24 hours, unless directed by a doctor
- children: take patches when told. administer to adults when told
 - if blood NCCC content falls below 0.40%, take 1-2 patches
 - directions apply irrespective of calendar age

Other Information

- read all warnings and directions before use. Keep carton.
- store between 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F) and excessive humidity

Made in USA

ORG 01/25

PRICE: USD \$6,000. CAN \$8,202.

Drug Facts (continued)

Inactive Ingredients

love, dedication, relaxation, comfort, pleasure, delight, gifts, remittances, phone calls, letters, heartfelt gestures, closeness, equality, humanity, freedom

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

EXP DATE: n/a

LOT:

†This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Tylenol®.

Distributed by:

Deflective Pharmaceuticals Corp.

Hauppauge, NY 11788